

# Bayesian Reasoning

## Chapter 13



[Thomas Bayes, 1701-1761](#)

# Today's topics

- Review probability theory
- Bayesian inference
  - From the joint distribution
  - Using independence/factoring
  - From sources of evidence

# Sources of Uncertainty

- Uncertain **inputs** -- missing and/or noisy data
- Uncertain **knowledge**
  - Multiple causes lead to multiple effects
  - Incomplete enumeration of conditions or effects
  - Incomplete knowledge of causality in the domain
  - Probabilistic/stochastic effects
- Uncertain **outputs**
  - Abduction and induction are inherently uncertain
  - Default reasoning, even deductive, is uncertain
  - Incomplete deductive inference may be uncertain
- ▶ Probabilistic reasoning only gives probabilistic results (summarizes uncertainty from various sources)

# Decision making with uncertainty

## **Rational** behavior:

- For each possible action, identify the possible outcomes
- Compute the **probability** of each outcome
- Compute the **utility** of each outcome
- Compute the probability-weighted (**expected**) **utility** over possible outcomes for each action
- Select action with the highest expected utility (principle of **Maximum Expected Utility**)

# Why probabilities anyway?

Kolmogorov showed that three simple axioms lead to the rules of probability theory

1. All probabilities are between 0 and 1:


$$0 \leq P(a) \leq 1$$

2. Valid propositions (tautologies) have probability 1, and unsatisfiable propositions have probability 0:

$$P(\text{true}) = 1 ; P(\text{false}) = 0$$

3. The probability of a disjunction is given by:

$$P(a \vee b) = P(a) + P(b) - P(a \wedge b)$$



# Probability theory 101

- **Random variables**
  - Domain
- **Atomic event:**  
complete specification of state
- **Prior probability:**  
degree of belief without any other evidence
- **Joint probability:**  
matrix of combined probabilities of a set of variables
- Alarm, Burglary, Earthquake
  - Boolean (like these), discrete, continuous
- $\text{Alarm}=\text{T} \wedge \text{Burglary}=\text{T} \wedge \text{Earthquake}=\text{F}$   
 $\text{alarm} \wedge \text{burglary} \wedge \neg \text{earthquake}$
- $P(\text{Burglary}) = 0.1$   
 $P(\text{Alarm}) = 0.1$   
 $P(\text{earthquake}) = 0.000003$
- $P(\text{Alarm, Burglary}) =$

	alarm	$\neg$ alarm
burglary	.09	.01
$\neg$ burglary	.1	.8

# Probability theory 101

	alarm	$\neg$ alarm
burglary	.09	.01
$\neg$ burglary	.1	.8

- **Conditional probability:** prob. of effect given causes
  - **Computing conditional probs:**
    - $P(a | b) = P(a \wedge b) / P(b)$
    - $P(b)$ : **normalizing** constant
  - **Product rule:**
    - $P(a \wedge b) = P(a | b) * P(b)$
  - **Marginalizing:**
    - $P(B) = \sum_a P(B, a)$
    - $P(B) = \sum_a P(B | a) P(a)$  (**conditioning**)
- $P(\text{burglary} | \text{alarm}) = .47$   
 $P(\text{alarm} | \text{burglary}) = .9$
  - $P(\text{burglary} | \text{alarm}) = P(\text{burglary} \wedge \text{alarm}) / P(\text{alarm}) = .09 / .19 = .47$
  - $P(\text{burglary} \wedge \text{alarm}) = P(\text{burglary} | \text{alarm}) * P(\text{alarm}) = .47 * .19 = .09$
  - $P(\text{alarm}) = P(\text{alarm} \wedge \text{burglary}) + P(\text{alarm} \wedge \neg \text{burglary}) = .09 + .1 = .19$

# Example: Inference from the joint

	alarm		¬alarm	
	earthquake	¬earthquake	earthquake	¬earthquake
burglary	.01	.08	.001	.009
¬burglary	.01	.09	.01	.79

$$\begin{aligned} P(\text{burglary} \mid \text{alarm}) &= \alpha P(\text{burglary}, \text{alarm}) \\ &= \alpha [P(\text{burglary}, \text{alarm}, \text{earthquake}) + P(\text{burglary}, \text{alarm}, \neg\text{earthquake})] \\ &= \alpha [ (.01, .01) + (.08, .09) ] \\ &= \alpha [ (.09, .1) ] \end{aligned}$$

Since  $P(\text{burglary} \mid \text{alarm}) + P(\neg\text{burglary} \mid \text{alarm}) = 1$ ,  $\alpha = 1/(\text{.09} + \text{.1}) = 5.26$   
(i.e.,  $P(\text{alarm}) = 1/\alpha = \text{.19}$  – **quizlet**: how can you verify this?)

$$P(\text{burglary} \mid \text{alarm}) = \text{.09} * 5.26 = \text{.474}$$

$$P(\neg\text{burglary} \mid \text{alarm}) = \text{.1} * 5.26 = \text{.526}$$

# Exercise:

## Inference from the joint



$p(\text{smart} \wedge \text{study} \wedge \text{prep})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

- **Queries:**
  - What is the prior probability of *smart*?
  - What is the prior probability of *study*?
  - What is the conditional probability of *prepared*, given *study* and *smart*?
- **Save these answers for next time! 😊**

# Independence



- When sets of variables don't affect each others' probabilities, we call them **independent**, and can easily compute their joint and conditional probability:  
Independent(A, B)  $\rightarrow$   $P(A \wedge B) = P(A) * P(B)$ ,  $P(A | B) = P(A)$
- {moonPhase, lightLevel} *might* be independent of {burglary, alarm, earthquake}
  - Maybe not: crooks may be more likely to burglarize houses during a new moon (and hence little light)
  - But if we know the light level, the moon phase doesn't affect whether we are burglarized
  - If burglarized, light level doesn't affect if alarm goes off
- Need a more complex notion of independence and methods for reasoning about the relationships



# Exercise: Independence

$p(\text{smart} \wedge \text{study} \wedge \text{prep})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

## Queries:

- Is *smart* independent of *study*?
- Is *prepared* independent of *study*?

# Conditional independence

- Absolute independence:
  - A and B are **independent** if  $P(A \wedge B) = P(A) * P(B)$ ;  
equivalently,  $P(A) = P(A | B)$  and  $P(B) = P(B | A)$
- A and B are **conditionally independent** given C if
  - $P(A \wedge B | C) = P(A | C) * P(B | C)$
- This lets us decompose the joint distribution:
  - $P(A \wedge B \wedge C) = P(A | C) * P(B | C) * P(C)$
- Moon-Phase and Burglary are *conditionally independent given* Light-Level
- Conditional independence is weaker than absolute independence, but still useful in decomposing the full joint probability distribution

# Exercise: Conditional independence




$p(\text{smart} \wedge \text{study} \wedge \text{prep})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

## Queries:

- Is *smart* conditionally independent of *prepared*, given *study*?
- Is *study* conditionally independent of *prepared*, given *smart*?

# Bayes' rule




- Derived from the product rule:
  - $P(C | E) = P(E | C) * P(C) / P(E)$
- Often useful for diagnosis:
  - If E are (observed) effects and C are (hidden) causes,
  - We may have a model for how causes lead to effects ( $P(E | C)$ )
  - We may also have prior beliefs (based on experience) about the frequency of occurrence of effects ( $P(C)$ )
  - Which allows us to reason abductively from effects to causes ( $P(C | E)$ )

# Ex: meningitis and stiff neck

- Meningitis (M) can cause a a stiff neck (S), though here are many other causes for S, too
- We'd like to use S as a diagnostic symptom and estimate  $p(M|S)$
- Studies can easily estimate  $p(M)$ ,  $p(S)$  and  $p(S|M)$   
 $p(M)=0.7$ ,  $p(S)=0.01$ ,  $p(S|M)=0.00002$
- Applying Bayes' Rule:  
$$p(M|S) = p(S|M) * p(M) / p(S) = 0.0014$$
- We can also do this w/o  $p(S)$  if we know  $p(S|\sim M)$   
 $\alpha \langle p(S|M)*P(m), p(S|\sim M)*p(\sim M) \rangle$

# Bayesian inference

- In the setting of diagnostic/evidential reasoning



- Know prior probability of hypothesis

conditional probability

$$P(H_i)$$

$$P(E_j | H_i)$$

$$P(H_i | E_j)$$

- Want to compute the *posterior probability*

- Bayes' s theorem (formula 1):

$$P(H_i | E_j) = P(H_i) * P(E_j | H_i) / P(E_j)$$

# Simple Bayesian diagnostic reasoning

- Also known as: Naive Bayes classifier
- Knowledge base:
  - Evidence / manifestations:  $E_1, \dots, E_m$
  - Hypotheses / disorders:  $H_1, \dots, H_n$ 
    - Note:  $E_j$  and  $H_i$  are **binary**; hypotheses are **mutually exclusive** (non-overlapping) and **exhaustive** (cover all possible cases)
  - Conditional probabilities:  $P(E_j | H_i), i = 1, \dots, n; j = 1, \dots, m$
- Cases (evidence for a particular instance):  $E_1, \dots, E_l$
- Goal: Find the hypothesis  $H_i$  with the highest posterior
  - $\text{Max}_i P(H_i | E_1, \dots, E_l)$

# Simple Bayesian diagnostic reasoning

- Bayes' rule says that

$$P(H_i | E_1 \dots E_m) = P(E_1 \dots E_m | H_i) P(H_i) / P(E_1 \dots E_m)$$

- Assume each evidence  $E_i$  is conditionally independent of the others, *given* a hypothesis  $H_i$ , then:

$$P(E_1 \dots E_m | H_i) = \prod_{j=1}^m P(E_j | H_i)$$

- If we only care about relative probabilities for the  $H_i$ , then we have:

$$P(H_i | E_1 \dots E_m) = \alpha P(H_i) \prod_{j=1}^m P(E_j | H_i)$$

# Limitations

- Cannot easily handle multi-fault situations, nor cases where intermediate (hidden) causes exist:
  - Disease D causes syndrome S, which causes correlated manifestations  $M_1$  and  $M_2$
- Consider a composite hypothesis  $H_1 \wedge H_2$ , where  $H_1$  and  $H_2$  are independent. What's the relative posterior?

$$\begin{aligned} P(H_1 \wedge H_2 \mid E_1, \dots, E_l) &= \alpha P(E_1, \dots, E_l \mid H_1 \wedge H_2) P(H_1 \wedge H_2) \\ &= \alpha P(E_1, \dots, E_l \mid H_1 \wedge H_2) P(H_1) P(H_2) \\ &= \alpha \prod_{j=1}^l P(E_j \mid H_1 \wedge H_2) P(H_1) P(H_2) \end{aligned}$$

- How do we compute  $P(E_j \mid H_1 \wedge H_2)$  ?

# Limitations

- Assume  $H_1$  and  $H_2$  are independent, given  $E_1, \dots, E_l$ ?
  - $P(H_1 \wedge H_2 \mid E_1, \dots, E_l) = P(H_1 \mid E_1, \dots, E_l) P(H_2 \mid E_1, \dots, E_l)$
- This is a very unreasonable assumption
  - Earthquake and Burglar are independent, but *not* given Alarm:
    - $P(\text{burglar} \mid \text{alarm}, \text{earthquake}) \ll P(\text{burglar} \mid \text{alarm})$
- Another limitation is that simple application of Bayes' s rule doesn't allow us to handle causal chaining:
  - A: this year's weather; B: cotton production; C: next year's cotton price
  - A influences C indirectly:  $A \rightarrow B \rightarrow C$
  - $P(C \mid B, A) = P(C \mid B)$
- Need a richer representation to model interacting hypotheses, conditional independence, and causal chaining
- Next: conditional independence and Bayesian networks!

# Summary

- Probability is a rigorous formalism for uncertain knowledge
- **Joint probability distribution** specifies probability of every **atomic event**
- Can answer queries by summing over atomic events
- But we must find a way to reduce the joint size for non-trivial domains
- **Bayes' rule** lets unknown probabilities be computed from known conditional probabilities, usually in the causal direction
- **Independence** and **conditional independence** provide the tools


# Postscript: Frequentists vs. Bayesians

- Frequentist inference draws conclusions from sample data based on the frequency or proportion of the data
- Bayesian inference uses Bayes' rule to update probability estimates for a hypothesis as additional evidence is learned
- The differences are often subtle, but can be consequential

DID THE SUN JUST EXPLODE?  
(IT'S NIGHT, SO WE'RE NOT SURE.)

# Frequentists vs. Bayesians

<http://xkcd.com/1132/>



FREQUENTIST STATISTICIAN:

THE PROBABILITY OF THIS RESULT HAPPENING BY CHANCE IS  $\frac{1}{36} = 0.027$ .  
SINCE  $p < 0.05$ , I CONCLUDE THAT THE SUN HAS EXPLODED.



BAYESIAN STATISTICIAN:

BET YOU \$50 IT HASN'T.

