# Bayesian Reasoning Chapters 12 & 13



Thomas Bayes, 1701-1761

# **Today's topics**

- Review probability theory
- Bayesian inference
  - -From the joint distribution
  - -Using independence/factoring
  - -From sources of evidence
- Naïve Bayes algorithm for inference and classification tasks

### **Many Sources of Uncertainty**

- Uncertain inputs -- missing and/or noisy data
- Uncertain knowledge
  - -Multiple causes lead to multiple effects
  - -Incomplete enumeration of conditions or effects
  - -Incomplete knowledge of causality in the domain
  - Probabilistic/stochastic effects
- Uncertain outputs
  - -Abduction and induction are inherently uncertain
  - Default reasoning, even deductive, is uncertain
  - -Incomplete deductive inference may be uncertain
  - Probabilistic reasoning only gives probabilistic results

#### **Decision making with uncertainty**

**Rational** behavior: for each possible action:

- Identify possible outcomes and for each
  - Compute **probability** of outcome
  - Compute **utility** of outcome
- Compute probability-weighted (expected) utility over possible outcomes
- Select action with the highest expected utility (principle of Maximum Expected Utility)

### Consider

- Your house has an alarm system
- It should go off if a burglar breaks into the house
- It can go off if there is an earthquake
- How can we predict what's happened if the alarm goes off?
  - -Someone has broken in!
  - -It's a minor earthquake



### **Probability theory 101**

#### • Random variables

– Domain

#### • Atomic event:

complete specification of state

#### • Prior probability:

degree of belief without any other evidence or info

#### Joint probability: matrix of combined probabilities of set of variables

- Alarm, Burglary, Earthquake Boolean (these), discrete (0-9), continuous (float)
- Alarm=T^Burglary=T^Earthquake=F alarm ^ burglary ^ -earthquake
- P(Burglary) = 0.1
   P(Alarm) = 0.1
   P(earthquake) = 0.000003
- P(Alarm, Burglary) =

	alarm	−alarm
burglary	.09	.01
¬burglary	.1	.8

### **Probability theory 101**

	alarm	−alarm
burglary	.09	.01
¬burglary	.1	.8

- Conditional probability: prob. of effect given causes
- Computing conditional probs:
  - $P(a | b) = P(a \land b) / P(b)$
  - P(b): normalizing constant
- Product rule:
  - − P(a ∧ b) = P(a | b) \* P(b)
- Marginalizing:
  - $P(B) = \Sigma_a P(B, a)$
  - $P(B) = \Sigma_a P(B | a) P(a)$ (conditioning)

- P(burglary | alarm) = .47
   P(alarm | burglary) = .9
- P(burglary | alarm) = P(burglary ^ alarm) / P(alarm) = .09/.19 = .47
- P(burglary \wedge alarm) =

   P(burglary | alarm) \* P(alarm)
   = .47 \* .19 = .09
- P(alarm) =  $P(alarm \land burglary) +$   $P(alarm \land \neg burglary)$ = .09+.1 = .19

#### **Example: Inference from the joint**

	ala	rm	−alarm	
	earthquake	¬earthquake	earthquake	
burglary	.01	.08	.001	.009
¬burglary	.01	.09	.01	.79

 $P(burglary | alarm) = \alpha P(burglary, alarm)$ 

=  $\alpha$  [P(burglary, alarm, earthquake) + P(burglary, alarm, ¬earthquake) =  $\alpha$  [ (.01, .01) + (.08, .09) ] =  $\alpha$  [ (.09, .1) ]

Since P(burglary | alarm) + P(¬burglary | alarm) = 1,  $\alpha = 1/(.09+.1) = 5.26$ (i.e., P(alarm) =  $1/\alpha = .19 - quizlet$ : how can you verify this?)

P(burglary | alarm) = .09 \* 5.26 = .474

 $P(\neg burglary | alarm) = .1 * 5.26 = .526$ 

### Consider



- A student has to take an exam
- She might be smart
- She might have studied
- She may be prepared for the exam
- How are these related?

#### smart -smart p(smart Λ study $\land$ prep) study -study -study study .16 prepared .432 .084 .008 .16 .036 .048 .072 -prepared

#### Queries:

- What is the prior probability of *smart*?
- What is the prior probability of study?
- What is the conditional probability of *prepared*, given study and smart?



#### **Exercise:**

#### Exercise: Inference from the joint

p(smart study ^ prep)	smart		smart	
	study	−study	study	−study
prepared	.432	.16	.084	.008
-prepared	.048	.16	.036	.072

#### Queries:

- What is the prior probability of *smart*?
- What is the prior probability of *study*?
- What is the conditional probability of *prepared*, given study and smart?

p(smart) = .432 + .16 + .048 + .16 = 0.8



p(smart ^ study ^ prep)	smart		smart	
	study	−study	study	−study
prepared	.432	.16	.084	.008
¬prepared	.048	.16	.036	.072

#### Queries:

- What is the prior probability of *smart*?
- What is the prior probability of *study*?
- What is the conditional probability of *prepared*, given study and smart?



#### **Exercise:**

p(smart 🔨	smart		smart	
study $\land$ prep)	study	−study	study	−study
prepared	.432	.16	.084	.008
-prepared	.048	.16	.036	.072

#### Queries:

**Exercise:** 

- What is the prior probability of *smart*?
- What is the prior probability of *study*?
- What is the conditional probability of *prepared*, given study and smart?

p(study) = .432 + .048 + .084 + .036 = **0.6** 



p(smart 🔨	smart		smart	
study $\land$ prep)	study	<b>¬study</b>	study	study
prepared	.432	.16	.084	.008
<b>¬prepared</b>	.048	.16	.036	.072

#### Queries:

**Exercise:** 

- What is the prior probability of *smart*?
- What is the prior probability of study?
- What is the conditional probability of *prepared*, given study and smart?



#### **Exercise:**

### Inference from the joint



p(smart 🔨	smart		smart	
study $\land$ prep)	study	_study	study	—study
prepared	.432	.16	.084	.008
prepared	.048	.16	.036	.072

#### Queries:

- What is the prior probability of *smart*?
- What is the prior probability of *study*?
- What is the conditional probability of *prepared*, given *study* and *smart*?

p(prepared | smart, study) = p(prepared, smart, study)/p(smart, study)= .432 / (.432 + .048) = 0.9

#### Independence



 When variables don't affect each others' probabilities, they are independent; we can easily compute their joint & conditional probability:

Independent(A, B)  $\rightarrow$  P(A $\land$ B) = P(A) \* P(B) or P(A|B) = P(A)

- {moonPhase, lightLevel} might be independent of {burglary, alarm, earthquake}
  - Maybe not: burglars may be more active during a new moon because darkness hides their activity
  - But if we know light level, moon phase doesn't affect whether we are burglarized
  - If burglarized, light level doesn't affect if alarm goes off
- Need a more complex notion of independence and methods for reasoning about the relationships



p(smart 🔨	smart		smart	
study $\land$ prep)	study	study	study	study
prepared	.432	.16	.084	.008
-prepared	.048	.16	.036	.072

#### **Queries:**

- -Q1: Is *smart* independent of *study*?
- -Q2: Is *prepared* independent of *study*?

How can we tell?



p(smart 🔨	smart		smart	
study $\land$ prep)	study	—study	study	—study
prepared	.432	.16	.084	.008
-prepared	.048	.16	.036	.072

#### Q1: Is *smart* independent of *study*?

- You might have some intuitive beliefs based on your experience
- You can also check the data

Which way to answer this is better?



p(smart $\wedge$	smart		smart	
study $\land$ prep)	study	_study	study	—study
prepared	.432	.16	.084	.008
-prepared	.048	.16	.036	.072

Q1: Is *smart* independent of *study*?

Q1 true iff p(smart|study) == p(smart)

**p(smart)** = .432 + 0.048 = .16 + .16 =

p(smart|study) = p(smart,study)/p(study)
= (.432 + .048) / .6 = 0.48/.6 = 0.8
0.8 == 0.8, so smart is independent of study



p(smart 🔨	smart		smart	
study $\land$ prep)	study	study	study	¬study
prepared	.432	.16	.084	.008
-prepared	.048	.16	.036	.072

#### Q2: Is *prepared* independent of *study*?

- What is prepared?
- •Q2 true iff



	p(smart ^ study ^ prep)	smart		smart	
		study	study	study	_study
	prepared	.432	.16	.084	.008
1	-prepared	.048	.16	.036	.072

#### Q2: Is *prepared* independent of *study*?

Q2 true iff p(prepared|study) == p(prepared) p(prepared) = .432 + .16 + .84 + .008 = .684 p(prepared|study) = p(prepared,study)/p(study) = (.432 + .084) / .6 = .86

 $0.86 \neq 0.684$ , so prepared not independent of study

#### **Absolute & conditional independence**

- Absolute independence:
  - A and B are **independent** if  $P(A \land B) = P(A) * P(B)$ ; equivalently, P(A) = P(A | B) and P(B) = P(B | A)
- A and B are conditionally independent given C if  $-P(A \land B | C) = P(A | C) * P(B | C)$
- This lets us decompose the joint distribution:

 $- P(A \land B \land C) = P(A | C) * P(B | C) * P(C)$ 

- Moon-Phase and Burglary are *conditionally independent given* Light-Level
- Conditional independence is weaker than absolute independence, but useful in decomposing full joint probability distribution

#### **Conditional independence**

- Intuitive understanding: conditional independence often comes from causal relations
  - -Moon phase causally effects light level at night
  - Other things do too, e.g., street lights
- For our burglary scenario, moon phase doesn't effect anything else
- Knowing *light level* means we can ignore moon phase in predicting whether or not alarm suggests we had a burglary

## **Bayes' rule**

Derived from the product rule:



-P(A, B) = P(A|B) \* P(B) # from definition of conditional probability

- -P(B, A) = P(B|A) \* P(A) # from definition of conditional probability
- -P(A, B) = P(B, A) # since order is not important

So...

P(A|B) = P(B|A) \* P(A)P(B)

# **Useful for diagnosis!**

- C is a cause, E is an effect: -P(C|E) = P(E|C) \* P(C) / P(E)
- Useful for diagnosis:
- E are (observed) effects and C are (hidden) causes,
- Often have model for how causes lead to effects P(E|C)
- May also have info (based on experience) on frequency of causes (P(C))
- Which allows us to reason <u>abductively</u> from effects to causes (P(C|E))



### Ex: meningitis and stiff neck

- Meningitis (M) can cause stiff neck (S), though there are other causes too
- Use S as a diagnostic symptom and estimate
   p(M|S)
- Studies can estimate p(M), p(S) & p(S|M), e.g. p(M)=0.7, p(S)=0.01, p(M)=0.00002
- Harder to directly gather data on p(M|S)
- Applying Bayes' Rule:
   p(M|S) = p(S|M) \* p(M) / p(S) = 0.0014

### Reasoning from evidence to a cause

• In the setting of diagnostic/evidential reasoning



hypotheses

evidence/manifestations

- Know prior probability of hypothesis $P(H_i)$ conditional probability $P(E_j | H_i)$
- Want to compute the *posterior probability*  $P(H_i | E_j)$
- Bayes' s theorem:

$$P(H_i | E_j) = P(H_i) * P(E_j | H_i) / P(E_j)$$

#### Simple Bayesian diagnostic reasoning

- Naive Bayes classifier
- Knowledge base:
  - Evidence / manifestations: E<sub>1</sub>, ... E<sub>m</sub>
  - Hypotheses / disorders: H<sub>1</sub>, ... H<sub>n</sub>

Note: E<sub>j</sub> and H<sub>i</sub> are **binary**; hypotheses are **mutually exclusive** (non-overlapping) and **exhaustive** (cover all possible cases)

- Conditional probabilities:  $P(E_j | H_i)$ , i = 1, ..., n; j = 1, ..., m
- Cases (evidence for a particular instance): E<sub>1</sub>, ..., E<sub>1</sub>
- Goal: Find the hypothesis  $H_i$  with highest posterior –  $Max_i P(H_i | E_1, ..., E_l)$

#### Simple Bayesian diagnostic reasoning

• Bayes' rule:

 $P(H_i | E_1...E_m) = P(E_1...E_m | H_i) P(H_i) / P(E_1...E_m)$ 

- Assume each evidence  $E_i$  is conditionally independent of the others, given a hypothesis  $H_i$ , then:  $P(E_1...E_m | H_i) = \prod_{j=1}^m P(E_j | H_j)$
- If only care about relative probabilities for  $H_i$ , then:  $P(H_i | E_1...E_m) = \alpha P(H_i) \prod_{j=1}^m P(E_j | H_j)$

### Limitations



- Can't easily handle **multi-fault situations** or cases where intermediate (hidden) causes exist:
  - Disease D causes syndrome S, which causes correlated manifestations M<sub>1</sub> and M<sub>2</sub>
- Consider composite hypothesis  $H_1 \wedge H_2$ , where  $H_1 \& H_2$  independent. What's relative posterior? P( $H_1 \wedge H_2 | E_1, ..., E_1$ ) =  $\alpha P(E_1, ..., E_1 | H_1 \wedge H_2) P(H_1 \wedge H_2)$ H<sub>2</sub>)
  - =  $\alpha P(E_1, ..., E_1 | H_1 \wedge H_2) P(H_1) P(H_2)$ =  $\alpha \prod_{j=1}^{l} P(E_j | H_1 \wedge H_2) P(H_1) P(H_2)$
- How do we compute  $P(E_j | H_1 \land H_2)$ ?

### Limitations



• Assume H1 and H2 independent, given E1, ..., El?

 $-P(H_1 \land H_2 | E_1, ..., E_l) = P(H_1 | E_1, ..., E_l) P(H_2 | E_1, ..., E_l)$ 

- Unreasonable assumption
  - Earthquake & Burglar independent, but not given Alarm:
     P(burglar | alarm, earthquake) << P(burglar | alarm)</li>
- Doesn't allow causal chaining:
  - A: 2017 weather; B: 2017 corn production; C: 2018 corn price
  - A influences C indirectly:  $A \rightarrow B \rightarrow C$
  - -P(C | B, A) = P(C | B)
- Need richer representation for interacting hypotheses, conditional independence & causal chaining
- Next: Bayesian Belief networks!

#### Summary



- Probability a rigorous formalism for uncertain knowledge
- Joint probability distribution specifies probability of every atomic event
- Answer queries by summing over atomic events
- Must reduce joint size for non-trivial domains
- Bayes rule: compute from known conditional probabilities, usually in causal direction
- Independence & conditional independence provide tools
- Next: Bayesian belief networks