

Levenshtein Distance Worksheet

(based on, and thanks to, Michael Gilletad <http://www.merriampark.com/ld.htm>)

Step	Description
1	Set n to be the length of s. Set m to be the length of t. If n = 0, return m and exit. If m = 0, return n and exit. Construct a matrix containing 0..m rows and 0..n columns.
2	Initialize the first row to 0..n. Initialize the first column to 0..m.
3	Examine each character of s (i from 1 to n).
4	Examine each character of t (j from 1 to m).
5	If s[i] equals t[j], the cost is 0. If s[i] doesn't equal t[j], the cost is 1. (and note in upper left of each cell)
6	Set cell d[i,j] of the matrix equal to the minimum of: a. The cell immediately above plus 1: d[i-1,j] + 1. b. The cell immediately to the left plus 1: d[i,j-1] + 1. c. The cell diagonally above and to the left plus the cost: d[i-1,j-1] + cost.
7	After the iteration steps (3, 4, 5, 6) are complete, the distance is found in cell d[n,m].

C A T

	0	1	2	3
R	1	1	2	3
A	2	2	0	1
T	3	3	2	0

P A R I S

	0	1	2	3	4	5	6
F	1	2	1	2	3	4	5
A	2	1	2	0	1	3	4
I	3	3	1	3	2	1	2
S	4	4	3	4	2	3	1
	5	4	5	3	3	2	3
	6						

Four ops to change Paris to fair: c/P/f, ins i,^R del i, del s