


# Bayesian Reasoning

Chapters 12 & 13



[Thomas Bayes, 1701-1761](#)

# Today's topics


- Motivation
- Review probability theory
- Bayesian inference
  - From the joint distribution
  - Using independence/factoring
  - From sources of evidence
- Naïve Bayes algorithm for inference and classification tasks

# Motivation: causal reasoning



- As the sun rises, the rooster crows
  - Does this correlation imply causality?
  - If so, which way does it go?
- The evidence can come from
  - Probabilities and Bayesian reasoning
  - Common sense knowledge
  - Experiments
- Bayesian Belief Networks (BBNs) are useful for modeling causal reasoning

# Motivation: logic isn't enough



- Classical logic is designed to work with certainties
- Getting a positive result on a COVID test doesn't necessarily mean you are infected
- And a negative result doesn't necessarily mean you are not infected
- You need to know the **true/false positive** and **true/false negative** rates of the test

# Decision making with uncertainty



**Rational** behavior: for each possible action:

- Identify possible outcomes and **for each**
  - Compute **probability** of outcome
  - Compute **utility** of outcome
  - Compute probability-weighted (**expected**) **utility** of outcome
- Select action with the highest expected utility (principle of **Maximum Expected Utility**)

# Consider

- Your house has an alarm system
- It should go off if a burglar breaks into the house
- It can also go off if there is an earthquake
- How can we predict what's happened if the alarm goes off?
  - Someone has broken in!
  - It's a minor earthquake



# Probability theory 101

- **Random variables:**

- Domain

- **Atomic event:**

- complete specification of state

- **Prior probability:**

- degree of belief without any other evidence or info

- **Joint probability:**

- matrix of combined probabilities of set of variables

- Alarm, Burglary, Earthquake

- Boolean (these) or discrete (0-9), continuous (float)

- Alarm=T $\wedge$ Burglary=T $\wedge$ Earthquake=F

- alarm  $\wedge$  burglary  $\wedge$   $\neg$ earthquake

- P(Burglary) = 0.1

- P(Alarm) = 0.1

- P(earthquake) = 0.000003

- P(Alarm, Burglary) =

	alarm	$\neg$ alarm
burglary	.09	.01
$\neg$ burglary	.1	.8

# Probability theory 101

	alarm	¬alarm
burglary	.09	.01
¬burglary	.1	.8

- **Conditional probability:** prob. of effect given causes
  - **Computing conditional probs:**
    - $P(a | b) = P(a \wedge b) / P(b)$
    - $P(b)$ : **normalizing** constant
  - **Product rule:**
    - $P(a \wedge b) = P(a | b) * P(b)$
  - **Marginalizing:**
    - $P(B) = \sum_a P(B, a)$
    - $P(B) = \sum_a P(B | a) P(a)$  (**conditioning**)
- $P(\text{burglary} | \text{alarm}) = .47$   
 $P(\text{alarm} | \text{burglary}) = .9$
  - $P(\text{burglary} | \text{alarm}) = P(\text{burglary} \wedge \text{alarm}) / P(\text{alarm}) = .09 / .19 = .47$
  - $P(\text{burglary} \wedge \text{alarm}) = P(\text{burglary} | \text{alarm}) * P(\text{alarm}) = .47 * .19 = .09$
  - $P(\text{alarm}) = P(\text{alarm} \wedge \text{burglary}) + P(\text{alarm} \wedge \neg\text{burglary}) = .09 + .1 = .19$



# Probability theory 101

	alarm	-alarm
burglary	.09	.01
-burglary	.1	.8

- **Conditional probability:** prob. of effect given causes

- **Computing conditional probs:**

- $P(a | b) = P(a \wedge b) / P(b)$

- $P(b)$ : **normalizing** constant

- **Product rule:**

- $P(a \wedge b) = P(a | b) * P(b)$

- **Marginalizing:**

- $P(B) = \sum_a P(B, a)$

- $P(B) = \sum_a P(B | a) P(a)$   
(**conditioning**)

- $P(\text{burglary} | \text{alarm}) = .47$   
 $P(\text{alarm} | \text{burglary}) = .9$

- $P(\text{burglary} | \text{alarm}) =$   
 $P(\text{burglary} \wedge \text{alarm}) / P(\text{alarm})$   
 $= .09 / .19 = .47$

- $P(\text{burglary} \wedge \text{alarm}) =$   
 $P(\text{burglary} | \text{alarm}) * P(\text{alarm})$   
 $= .47 * .19 = .09$

- $P(\text{alarm}) =$   
 $P(\text{alarm} \wedge \text{burglary}) +$   
 $P(\text{alarm} \wedge \neg \text{burglary})$   
 $= .09 + .1 = .19$

# Example: Inference from the joint

	alarm		¬alarm	
	earthquake	¬earthquake	earthquake	¬earthquake
burglary	.01	.08	.001	.009
¬burglary	.01	.09	.01	.79

$$\begin{aligned} P(\text{burglary} \mid \text{alarm}) &= \alpha P(\text{burglary}, \text{alarm}) \\ &= \alpha [P(\text{burglary}, \text{alarm}, \text{earthquake}) + P(\text{burglary}, \text{alarm}, \neg\text{earthquake})] \\ &= \alpha [ (.01, .01) + (.08, .09) ] \\ &= \alpha [ (.09, .1) ] \end{aligned}$$

Since  $P(\text{burglary} \mid \text{alarm}) + P(\neg\text{burglary} \mid \text{alarm}) = 1$ ,  $\alpha = 1/(\text{.09} + \text{.1}) = 5.26$   
(i.e.,  $P(\text{alarm}) = 1/\alpha = \text{.19}$  – **quizlet**: how can you verify this?)

$$P(\text{burglary} \mid \text{alarm}) = \text{.09} * 5.26 = \text{.474}$$

$$P(\neg\text{burglary} \mid \text{alarm}) = \text{.1} * 5.26 = \text{.526}$$

# Consider

- A student has to take an exam
  - She might **be smart**
  - She might **have studied**
  - She may **be prepared** for the exam
- How are these related?
- We can collect joint probabilities for the three events
  - Measure “prepared” as “got a passing grade”



# Exercise: Inference from the joint



$p(\text{smart} \wedge \text{study} \wedge \text{prepared})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

Each of the 8 highlighted boxes has the joint probability for the three values of smart, study, prepared

## Queries:

- What is the prior probability of *smart*?
- What is the prior probability of *study*?
- What is the conditional probability of *prepared*, given *study* and *smart*?

Standard way to show joint probabilities of 3 variables as a 2D table

# Exercise:

## Inference from the joint



$p(\text{smart} \wedge \text{study} \wedge \text{prepared})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

### Queries:

- What is the prior probability of *smart*?
- What is the prior probability of *study*?
- What is the conditional probability of *prepared*, given *study* and *smart*?

$$p(\text{smart}) = .432 + .16 + .048 + .16 = \mathbf{0.8}$$

# Exercise:

## Inference from the joint



$p(\text{smart} \wedge \text{study} \wedge \text{prepared})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

### Queries:

- What is the prior probability of *smart*?
- **What is the prior probability of *study*?**
- What is the conditional probability of *prepared*, given *study* and *smart*?

# Exercise:

## Inference from the joint



$p(\text{smart} \wedge \text{study} \wedge \text{prepared})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

### Queries:

- What is the prior probability of *smart*?
- **What is the prior probability of *study*?**
- What is the conditional probability of *prepared*, given *study* and *smart*?

$$p(\text{study}) = .432 + .048 + .084 + .036 = 0.6$$

# Exercise:

## Inference from the joint



$p(\text{smart} \wedge \text{study} \wedge \text{prepared})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

### Queries:

- What is the prior probability of *smart*?
- What is the prior probability of *study*?
- **What is the conditional probability of *prepared*, given *study* and *smart*?**



# Exercise:

## Inference from the joint



$p(\text{smart} \wedge \text{study} \wedge \text{prepared})$	smart		$\neg\text{smart}$	
	study	$\neg\text{study}$	study	$\neg\text{study}$
prepared	.432	.16	.084	.008
$\neg\text{prepared}$	.048	.16	.036	.072

### Queries:

- What is the prior probability of *smart*?
- What is the prior probability of *study*?
- **What is the conditional probability of *prepared*, given *study* and *smart*?**

$$\begin{aligned} p(\text{prepared} | \text{smart}, \text{study}) &= p(\text{prepared}, \text{smart}, \text{study}) / p(\text{smart}, \text{study}) \\ &= .432 / (.432 + .048) \\ &= \mathbf{0.9} \end{aligned}$$

# Independence



- When variables don't affect each others' probabilities, they are **independent**; we can easily compute their joint & conditional probability:

$$\text{Independent}(A, B) \rightarrow P(A \wedge B) = P(A) * P(B); P(A | B) = P(A)$$

- {moonPhase, lightLevel} *might* be independent of {burglary, alarm, earthquake}
  - Maybe not: burglars may be more active during a new moon because darkness hides their activity
  - But if we know light level, moon phase doesn't affect whether we are burglarized
  - If burglarized, light level doesn't affect if alarm goes off
- Need a more complex notion of independence and methods for **reasoning about the relationships**



# Exercise: Independence

$p(\text{smart} \wedge \text{study} \wedge \text{prepared})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

## Queries:

- Q1: Is *smart* independent of *study*?
- Q2: Is *prepared* independent of *study*?

How can we tell?



# Exercise: Independence

$p(\text{smart} \wedge \text{study} \wedge \text{prepared})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

**Q1: Is *smart* independent of *study*?**

- You might have some **intuitive beliefs** based on your experience
- You can also **check the data**

Which way to answer this is better?

# Exercise: Independence



$p(\text{smart} \wedge \text{study} \wedge \text{prepared})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

**Q1: Is *smart* independent of *study*?**

**Q1 true iff  $p(\text{smart} | \text{study}) == p(\text{smart})$**

$$p(\text{smart}) = .432 + 0.048 + .16 + .16 = \mathbf{0.8}$$

$$p(\text{smart} | \text{study}) = p(\text{smart}, \text{study}) / p(\text{study}) \\ = (.432 + .048) / .6 = 0.48 / .6 = \mathbf{0.8}$$

**$0.8 == 0.8 \therefore \text{smart is independent of study}$**



# Exercise: Independence

$p(\text{smart} \wedge \text{study} \wedge \text{prep})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

**Q2: Is *prepared* independent of *study*?**

- What is prepared?
- Q2 true iff



# Exercise: Independence

$p(\text{smart} \wedge \text{study} \wedge \text{prep})$	smart		$\neg$ smart	
	study	$\neg$ study	study	$\neg$ study
prepared	.432	.16	.084	.008
$\neg$ prepared	.048	.16	.036	.072

**Q2: Is *prepared* independent of *study*?**

**Q2 true iff  $p(\text{prepared} | \text{study}) == p(\text{prepared})$**

$$p(\text{prepared}) = .432 + .16 + .084 + .008 = .684$$

$$p(\text{prepared} | \text{study}) = p(\text{prepared}, \text{study}) / p(\text{study})$$

$$= (.432 + .084) / .6 = .86$$

**$0.86 \neq 0.684, \therefore$  prepared not independent of study**

# Absolute & conditional independence

- Absolute independence:
  - A and B are **independent** if  $P(A \wedge B) = P(A) * P(B)$ ;  
equivalently,  $P(A) = P(A | B)$  and  $P(B) = P(B | A)$
- A and B are **conditionally independent** given C if
  - $P(A \wedge B | C) = P(A | C) * P(B | C)$


If it holds, lets us decompose the joint distribution:

- $P(A \wedge B \wedge C) = P(A | C) * P(B | C) * P(C)$
- Moon-Phase and Burglary are ***conditionally independent given*** Light-Level
- Conditional independence is weaker than absolute independence, but useful in decomposing full joint probability distribution




# Conditional independence

- Conditional independence often comes from **causal relations**
  - FullMoon causally affects LightLevel at night as does StreetLights
- In burglary scenario, FullMoon doesn't affect anything else
- Knowing *LightLevel*, we can ignore *FullMoon* and *StreetLights* when predicting if alarm suggests **Burglary**



# Bayes' rule



Derived from the product rule:

- $P(A, B) = P(A | B) * P(B)$  *# from definition of conditional probability*
- $P(B, A) = P(B | A) * P(A)$  *# from definition of conditional probability*
- $P(A, B) = P(B, A)$  *# since order is not important*

So...

$$P(A | B) = \frac{P(B | A) * P(A)}{P(B)}$$

relates  $P(A|B)$   
and  $P(B|A)$

$P(A,B)$  is probability of both A and B being true, so  $P(A,B) = P(B,A)$


# Useful for diagnosis!

- *C is a cause, E is an effect:*

- $P(C|E) = P(E|C) * P(C) / P(E)$

- **Useful for diagnosis:**

- E are (observed) effects and C are (hidden) causes,
  - Often have model for how causes lead to effects  $P(E|C)$
  - We may have info (based on experience) on frequency of causes ( $P(C)$ )
  - Which allows us to reason abductively from effects to causes ( $P(C|E)$ )
  - Recall, abductive reasoning: from  $A \Rightarrow B$  and  $B$ , infer (maybe?)  $A$



# Example: meningitis and stiff neck

cause

symptom

- **Meningitis (M)** can cause **stiff neck (S)**, though there are other causes too
- Use *S* as a *diagnostic symptom* & estimate  **$p(M|S)$**
- Studies can estimate  $p(M)$ ,  $p(S)$  &  $p(S|M)$ , e.g.  
 $p(S|M)=0.7$ ,  $p(S)=0.01$ ,  $p(M)=0.00002$
- Harder to directly gather data on  $p(M|S)$
- Applying Bayes' Rule:  
$$p(M|S) = p(S|M) * p(M) / p(S) = 0.0014$$



# Summary

- Probability a rigorous formalism for uncertain knowledge
- **Joint probability distribution** specifies probability of every **atomic event**
- Answer queries by summing over atomic events
- Must reduce joint size for non-trivial domains
- **Bayes rule**: compute from known conditional probabilities, usually in causal direction
- **Independence & conditional independence** provide tools
- Next: Bayesian belief networks